

A resolution of quantum dynamical semigroups

Anilesh Mohari

S.N.Bose Center for Basic Sciences,

JD Block, Sector-3, Calcutta-91

E-mail:anilesh@boson.bose.res.in

Abstract

We consider a class of quantum dissipative systems governed by a one parameter completely positive maps on a von-Neumann algebra. We introduce a notion of recurrent and metastable projections for the dynamics and prove that the unit operator can be decomposed into orthogonal projections where each projections are recurrent or metastable for the dynamics.

1 Introduction:

Let $\tau = (\tau_t, t \geq 0)$ be a semigroup of identity preserving completely positive maps [Da1,Da2,BR] on a von-Neumann algebra $\mathcal{A}_0$ acting on a separable Hilbert space $\mathcal{H}_0$, where either the parameter $t \in R_+$, the set of positive real numbers or $t \in Z^+$, the set of positive integers. In case $\mathcal{A}_0 = l^\infty(\mathcal{S})$, the bounded measurable functions on a countable set $\mathcal{S}$, τ is the Markov semigroup associated with a continuous or discrete time Markov chain. In such a case the state space $\mathcal{S}$ is well classified as disjoint union of recurrent and metastable states. Such a classification enriched our understanding of asymptotic behavior of associated stochastic processes. In this exposition we aim to achieve a similar classification in the general framework of non-commutative probability and prove results related to the asymptotic behavior of the quantum dynamical systems (τ_t) if it admits a normal invariant state. To that end we assume further that the map τ_t is normal for each $t \geq 0$ and the map $t \rightarrow \tau_t(x)$ is weak* continuous for each $x \in \mathcal{A}_0$.

Following [FR3] we say a projection $p \in \mathcal{A}_0$ is *sub-harmonic and harmonic* if $\tau_t(p) \geq p$ and $\tau_t(p) = p$ for all $t \geq 0$ respectively. For a sub-harmonic projection p , we define the reduced quantum dynamical semigroup (τ_t^p) on the von-Neumann algebra $p\mathcal{A}_0p$ by $\tau_t^p(x) = p\tau_t(x)p$ where $t \geq 0$ and $x \in \mathcal{A}_0^p$. If p is also a harmonic projection, the reduced dynamics τ_t^{1-p} and τ_t^p are complement to each other. For more details we refer to [Ev]. Following Evans [Ev] we also say (τ_t) is *irreducible* if there exists no non-trivial harmonic projection in $\mathcal{A}_0$ for (τ_t) . This notion is not quite equivalent to the notion of irreducibility for Markov chain [No].

A non-zero sub-harmonic projection p is said to be *minimal* if there exists

no sub-harmonic projection q so that $0 < q < p$. A non-zero projection $p \in \mathcal{A}$ is called *recurrent* if p is minimal sub-harmonic. A recurrent projection p is called *positive recurrent* if there exists a normal invariant state for (τ_t) with support equal to p . A recurrent projection is called *null recurrent* if it is not a positive recurrent. If the identity operator is a recurrent projection, we say (τ_t) is recurrent.

For a sub-harmonic projection p , 1 is an upper bound for the increasing positive operators $\tau_t(p)$, $t \geq 0$. Thus there exists an operator $0 \leq y \leq 1$ so that $y = \text{s.lim}_{t \rightarrow \infty} \tau_t(p)$. If y is an injective map on the Hilbert space $\mathcal{H}_0$ we say the projection $1 - p$ is *metastable*. If $y = 1$ we say $1 - p$ is *transient*. A normal state ϕ_0 is called *invariant* for (τ_t) if $\phi_0 \tau_t(x) = \phi_0(x)$ for all $x \in \mathcal{A}_0$ and $t \geq 0$. The support p of a normal invariant state is a sub-harmonic projection [FR3] and ϕ_0^p , the restriction of ϕ_0 to $\mathcal{A}_0^p$ is an invariant normal state for (τ_t^p) . Thus asymptotic properties (ergodic, mixing) of the dynamics $(\mathcal{A}_0, \tau_t, \phi_0)$ is well determined by the asymptotic properties (ergodic, mixing respectively) of the reduced dynamics $(\mathcal{A}_0^p, \tau_t^p, \phi_0^p)$ provided $y = 1$. For more details we refer to [Fr,Mol].

We explore a necessary and sufficient condition for a sub-harmonic projection to be metastable and also find a sufficient condition for a metastable projection to be transient. In case the von-Neumann algebra is type-I with centre completely atomic, we will show that there exists a mutually orthogonal family of recurrent projections (p_i) so that $\sum_i p_i \leq 1$ and $1 - \sum_i p_i$ is a metastable projection. However such a decomposition is not unique in general unlike in the classical Markov chain. Thus the problem remain open also for more general non-commutative algebra of observables. Related results for stochastically complete Brownian flows on a smooth manifold associated with

a family of vector fields are proved in [Mo2].

In particular for any finite dimensional Hilbert space $\mathcal{H}_0$ we prove that any metastable projection is also transient and any recurrent projection is also positive recurrent. Thus even in these general non-commutative dynamical systems of irreversible processes classical feature remain valid once the Hilbert space is finite dimensional. If there exists a unique recurrent projection p and $\mathcal{H}_0$ is finite dimensional then there exists an unique invariant normal state and (τ_t) is ergodic with respect to that state.

2 Sub-harmonic projections:

In the following we list few crucial property (Proposition 2.1 and Proposition 2.2) of sub-harmonic projections appeared in [FR3,Mo1].

PROPOSITION 2.1: Let p be a sub-harmonic projection for (τ_t) . Then the following hold:

- (a) for all $t \geq 0$, $p\tau_t(p) = \tau_t(p)p = p$.
- (b) $\tau_t(x(1-p))p = 0$ for all $x \in \mathcal{A}_0, t \geq 0$.

For a projection p , $\mathcal{A}_0^p = p\mathcal{A}_0p$ is also a von-Neumann algebra acting on the Hilbert subspace $p\mathcal{H}_0$. Thus for a sub-harmonic projection p we verify by Proposition 2.1 that (τ_t^p) defined by $\tau_t^p(x) = p\tau_t(x)p$, $x \in \mathcal{A}_0^p$ is a Markov semigroup. Let the strong limit of $\tau_t(p)$ be y as $t \uparrow \infty$. By Proposition 2.1 (a) we have $py = yp = p, p \leq y \leq 1$ and $\tau_t(y) = y \forall t \geq 0$. Thus $p\tau_t(1-y^2)p = p\tau_t(p(1-y^2)p)p = 0$. So we also have $p\tau_t(y^2) = \tau_t(y^2)p = p$ for all $t \geq 0$. Since $\tau_t(p) \leq \tau_t(y^2) \leq \tau_t(y) = y$, the strong limit of $\tau_t(y^2)$ as $t \rightarrow \infty$ is also y . In case y^2 is also an invariant element for (τ_t) , we have $y^2 = y$. In general y^2 need not be an invariant element even for an irreducible classical Markov semigroup

(τ_t) . For more details we refer to [Mo1].

PROPOSITION 2.2: Let p be a sub-normal projection and $y = \text{s.lim} \tau_t(p)$. Then for any $z \in \mathcal{B}(\mathcal{H}_0)$ the following statements are equivalent:

- (a) $yz = 0$
- (b) $\tau_t(p)z = 0$ for all $t \geq 0$.

The following proposition is crucial for classification and related problems.

PROPOSITION 2.3: Let $x \in \mathcal{A}_0$ be a non-negative invariant element for a completely positive unital normal map τ on a von-Neumann algebra $\mathcal{A}_0$ and q be the projection on the closure of the range of x . Then $1 - q$ is a sub-harmonic projection for τ .

PROOF: Let q be the unique minimal projection so that $(1 - q)x = 0$. For any unitary element $u' \in \mathcal{A}'_0$ in the commutant, we have $(1 - q)u'xu'^* = 0$, thus $u'^*(1 - q)u'x = 0$. Since $u'^*(1 - q)u'$ is also a projection, by minimality we conclude that $u'^*qu' \geq q$. u' being an arbitrary unitary element in $\mathcal{A}'_0$, we conclude that $u'^*qu' = q$. That q is an element in $\mathcal{A}_0$ follows now by von-Neumann's density theorem [BR]. By Steinspring representation [BR], there exists a countable family of bounded operators (l_k) so that $\tau(z) = \sum_k l_k^* z l_k$ for all $z \in \mathcal{A}_0$. Since x is an invariant element, we have $(1 - q)\tau(x) = (1 - q)x = 0$. Hence we have by the representation $xl_k(1 - q) = 0$. Thus we have $ql_k(1 - q) = 0$. Once more by the representation we conclude that $\tau(q)(1 - q) = 0$, i.e. $\tau(q) \leq q$. ■

For the classification we need to guarantee existence of a minimal projection.

THEOREM 2.4: Let $\mathcal{A}_0$ be type-I with centre completely atomic. There exists a commuting family (p_i) of countable many orthogonal recurrent projec-

tions for (τ_t) so that $q = 1 - \sum_i p_i$ is metastable.

PROOF: Since $\mathcal{A}_0$ is type-I with centre completely atomic, minimal projection exists [BR,vol-1]. In case there exists no non-trivial minimal sub-harmonic projection, we have trivial partition $p_1 = 1$ and $q = 0$. Suppose p_1 is a minimal sub-harmonic projection and q_1 is the projection on the closure of the range of y_1 where y_1 is the strong limit of $\tau_t(p_1)$ as before. Thus by Proposition 2.3 $1 - q_1$ is sub-harmonic. In case $1 - q_1 = 0$ or $1 - q_1$ is recurrent then $1 - p_1 - p_2$ is metastable, where $p_2 = 1 - q_1$. So the construction ends. Otherwise we look for a minimal sub-harmonic projection $0 < p_2 < (1 - q_1)$ and so on. Note that by our construction the family is countable ($\mathcal{H}_0$ is separable) and commuting.

■

THEOREM 2.5: Let p be a sub-harmonic projections for (τ_t) so that $y \geq \lambda$ for some $\lambda > 0$. Then $y = 1$.

PROOF: Since $yp = p$, by our hypothesis we write $y \geq p + \lambda(1 - p)$ for some $\lambda > 0$. Thus for any $t \geq t_0$ we have $y \geq \tau_t(p) + \lambda(1 - \tau_t(p))$. By taking limit $t \rightarrow \infty$ we conclude that $y \geq y + \lambda(1 - y)$. Hence $y \geq 1$, since $\lambda > 0$. Since $0 \leq y \leq 1$ we conclude that $y = 1$.

■

COROLLARY 2.6: Let p be a maximal non-trivial sub-harmonic projection $p \neq 1$ for (τ_t) and $1 - p$ be finite dimensional. Then $\tau_t(p) \rightarrow 1$ as $t \rightarrow \infty$ in strong operator topology.

PROOF: Let q be the projection on the range of y . Since y is a non-negative invariant element for (τ_t) , Proposition 2.3 says that $1 - q$ is sub-harmonic. Since p is a maximal non-trivial sub-harmonic projection which commutes with $1 - q$, we have $1 - q = 0$. Hence y is one to one. Since $1 - p$ is finite dimensional, y is bounded away from zero i.e. there exists a constant $\lambda > 0$ so that $y \geq \lambda 1$. Thus the proof is complete once we appeal to Theorem 2.5.

■

If the parameter $t \in Z_+$, the dynamics is determined by a unital completely positive normal map (τ) and $\tau_n = \tau \circ \dots \circ \tau$ - n fold composition. In such a case p is sub-harmonic for (τ_n) if and only if $\tau(p) \geq p$.

THEOREM 2.7: Let τ be a unital completely positive map on a von-Neumann algebra $\mathcal{A}_0$. Then p be a sub-harmonic projection for τ if and only if $(1-p)l_k p = 0$ for all $k \geq 1$, where (l_k) is the family of bounded operators so that $\tau(x) = \sum_k l_k^* x l_k$ for all $x \in \mathcal{A}_0$.

Moreover for a sub-harmonic projection p for τ , and a bounded operator z , $yz = 0$ if and only if $p l_{i_1} l_{i_2} l_{i_3} \dots l_{i_n} z = 0$, where $s.\lim_{n \rightarrow \infty} \tau_n(p) = y$.

PROOF: Let p be sub-harmonic for τ , i.e. $\tau(p) \geq p$. We have $p\tau(p)p \geq p$, p being a projection. Hence $p\tau(1-p)p \leq 0$. Since $1-p \geq 0$, we also have $p\tau(1-p)p \geq 0$. So $p\tau(1-p)p = 0$. By the representation we have $\sum_k p l_k^* (1-p) l_k p = 0$, hence the result follows. The last part follows once we note that $yz = 0$ if and only if $\tau_n(p)z = 0$ for all $n \geq 0$. ■

Hence y is injective if and only if p together with the range of these $l_{i_1}^* l_{i_2}^* \dots l_{i_m}^* p$ family of operators, where $i_1, i_2, \dots, i_m \geq 1$ generate the Hilbert space $\mathcal{H}_0$. In such a case for finite dimensional Hilbert space $\mathcal{H}_0$, Theorem 2.5 says that $y = 1$. Although the family (l_k) is not uniquely determined by the map τ , the criteria appeared in Theorem 2.7 is independent of the choice we make for it's representation.

3 Strong ergodic property:

In the last section we investigated various criteria for the support projection p associated with the normal invariant state so that $1-p$ is transient for the dynamics $(\mathcal{A}_0, \tau_t, \phi_0)$. In a recent paper [Mo1], we have investigated how various

asymptotic properties (ergodic, weak mixing, strong mixing) of $(\mathcal{A}_0, \tau_t, \phi_0)$ can be determined by the reduced dynamics $(\mathcal{A}_0^p, \tau_t^p, \phi_0)$ when $1 - p$ is transient. In the following theorem we prove one more such a property known in the literature as strong ergodicity.

THEOREM 3.1: Let $(\mathcal{A}_0, \tau_t, \phi_0)$ be a quantum dynamical systems with a normal invariant state ϕ_0 with support projection p . If $\text{s-limit}_{t \rightarrow \infty} \tau_t(p) = 1$ then the following statements are equivalent:

- (a) $\|\phi\tau_t - \phi_0\| \rightarrow 0$ as $t \rightarrow \infty$ for any normal state on $\mathcal{A}_0$.
- (b) $\|\phi\tau_t^p - \phi_0\| \rightarrow 0$ as $t \rightarrow \infty$ for any normal state on $\mathcal{A}_0$.

PROOF: That (b) implies (a) is trivial. For the converse we write $\|\phi\tau_t - \phi_0\| = \sup_{x: \|x\| \leq 1} |\phi\tau_t(x) - \phi_0(x)| \leq \sup_{\{x: \|x\| \leq 1\}} |\phi\tau_t(pxp) - \phi_0(pxp)| + \sup_{\{x: \|x\| \leq 1\}} |\phi\tau_t(pp^\perp x)| + \sup_{\{x: \|x\| \leq 1\}} |\phi\tau_t(p^\perp xp)| + \sup_{\{x: \|x\| \leq 1\}} |\phi\tau_t(p^\perp xp^\perp)|$. Since $\tau_t((1 - p)x) \rightarrow 0$ in the weak* topology and $|\phi\tau_t(xp^\perp)|^2 \leq |\phi\tau_t(xx^*)| |\phi(\tau_t(p^\perp))| \leq \|x\|^2 \phi(\tau_t(p^\perp))$ it is good enough if we verify that (a) is equivalent to $\sup_{\{x: \|x\| \leq 1\}} |\phi\tau_t(pxp) - \phi_0(pxp)| \rightarrow 0$ as $t \rightarrow \infty$. To that end we first note that $\limsup_{t \rightarrow \infty} \sup_{x: \|x\| \leq 1} |\psi(\tau_{s+t}(pxp)) - \phi_0(pxp)|$ is independent of $s \geq 0$ we choose. On the other hand we write $\tau_{s+t}(pxp) = \tau_s(p\tau_t(pxp)p) + \tau_s(p\tau_t(pxp)p^\perp) + \tau_s(p^\perp \tau_t(pxp)p) + \tau_s(p^\perp \tau_t(pxp)p^\perp)$ and use the fact for any normal state ϕ we have $\limsup_{t \rightarrow \infty} \sup_{x: \|x\| \leq 1} |\psi(\tau_s(z\tau_t(pxp)p^\perp))| \leq \|z\| |\psi(\tau_s(p^\perp))|$ for all $z \in \mathcal{A}_0$. Thus by our hypothesis on the support and we conclude that (a) hold whenever (b) is true. ■

4 Quantum mechanical master equation:

In this section we aim to deal with a class of quantum dynamical semigroup. We say a normal Markov semigroup (τ_t) on $\mathcal{A}_0$ is norm continuous if $\lim_{t \rightarrow 0} \|\tau_t -$

$I|| = 0$. In such a case the generator $\mathcal{L}$ is a bounded operator on $\mathcal{A}$ and can be described [GKS,Li,CE] by

$$\mathcal{L}(x) = Y^*x + xY + \sum_{k \geq 1} L_k^*xL_k \quad (4.1)$$

where $Y \in \mathcal{A}_0$ is the generator of a norm continuous contractive semigroup on $\mathcal{H}_0$ and L_k , $k \geq 1$ is a family of bounded operators so that $\sum_k L_k^*xL_k \in \mathcal{A}_0$ whenever $x \in \mathcal{A}_0$. However this choice $(Y, L_k, k \geq 1)$ is not unique. Conversely, for any such a family (Y, L_k) with $Y \in \mathcal{A}_0$ and $L_kxL_k^* \in \mathcal{A}_0$, $\forall x \in \mathcal{A}_0$, there exists a unique Markov semigroup (τ_t) with $\mathcal{L}$ as its generator. There are many methods to show the existence of a Markov semigroup (τ_t) with $\mathcal{L}$ as it's generator [Da3,MS,CF]. Here we describe one such a method [CF].

We consider the following iterated equation :

$$\tau_t^0(x) = e^{tY^*}xe^{tY} \quad (4.2)$$

$$\tau_t^{(n)}(x) = e^{tY^*}xe^{tY} + \int_0^t e^{(t-s)Y^*}\Phi(\tau_s^{(n-1)}(x))e^{(t-s)Y}ds, \quad n \geq 1 \quad (4.3)$$

where $\Phi(x) = \sum_k L_k^*xL_k$. It is simple to check for $x \geq 0$ that

$$0 \leq \tau_t^{n-1}(x) \leq \tau_t^n(x) \leq ||x|| I, \quad \forall t \geq 0$$

Thus we set for $x \geq 0$, $\tau_t(x) = \lim_{n \rightarrow \infty} \tau_t^{(n)}(x)$ in the weak* topology. For an arbitrary element we extend it by linearity. Thus we have

$$\tau_t(x) = e^{tY^*}xe^{tY} + \int_0^t e^{(t-s)Y^*}\Phi(\tau_s(x))e^{(t-s)Y}ds \quad (4.4)$$

for any $x \in \mathcal{A}_0$.

In such a case [Ev], it is simple to check that (τ_t) is irreducible if and only if $\{x \in \mathcal{A}_0 : [x, H] = 0, [x, L_k] = 0 \forall k \geq 1\}$ is trivial.

The following simple but important result due to Fagnola-Rebolledo [FR3].

THEOREM 4.1: A projection p is sub-harmonic if and only if $(1-p)Yp = 0$ and $(1-p)L_k p = 0$ for all $1 \leq k \leq \infty$.

PROOF: For a proof and a more general result we refer to [FR3]. ■

PROPOSITION 4.2: Let p be a sub-harmonic projection and $y = s.\lim_{t \rightarrow \infty} \tau_t(p)$. For any $z \in \mathcal{B}(\mathcal{H}_0)$ following are equivalent:

- (a) $yz = 0$
- (b) $pz = 0$, $pL_{i_1}L_{i_2}....L_{i_n}z = 0$ for all $0 \leq i_m < \infty$, $1 \leq m \leq n$ and $n \geq 1$, where $L_0 = Y$.

PROOF: By Proposition 2.2, $yz = 0$ if and only if $z^*\tau_t(p)z = 0$ for all $t \geq 0$. Now by (4.4) we have $z^*\tau_t(p)z = 0$ if and only if $z^*e^{tY^*}pe^{tY}z = 0$ and $z^*\Phi(\tau_t(p))z = 0$ for $t \geq 0$. Thus we have $pe^{tY}z = 0$ and also $z^*\Phi(\tau_t(p))z = 0$ for all t . Thus in particular we have $z^*\Phi(p)z = 0$, hence $pL_k z = 0$ for all $k \geq 0$. We go now by induction on n , we check if $z' = pL_{i_1}L_{i_2}....L_{i_n}z$ then $yz' = z'$ thus (a) implies (b). For the converse statement, we check that derivative of any order at $t = 0$ of $z^*\tau_t(p)z$ vanishes, thus constant which is zero. ■

Thus the zero operator is the only element z that satisfies (b) if and only if the closure of the range of y is the entire Hilbert space. Thus p together with $L_{i_1}^*L_{i_2}^*....L_{i_n}^*p$ where $0 \leq i_m < \infty$, $1 \leq m \leq n$ and $n \geq 1$ will generate the Hilbert space if and only if y is one to one. In particular we find this property is a necessary condition for y to be 1. In general the condition is not a sufficient one. Once more one can construct a counter example in birth and death processes where no population is an absorbing state and birth and death rates are such that the population will extinct with positive probability but need not be 1. We omit the details. However Corollary 2.6 says that for finite

dimensional $1 - p$ the condition is also sufficient for $y = 1$. So we have the following result.

THEOREM 4.3: Let p be a sub-harmonic projection for (τ_t) and $1 - p$ be finite dimensional. Then $y = 1$ if and only if p together with the ranges of operators $L_{i_1}^* \dots L_{i_n}^* p$, $i_1, \dots, i_n \geq 0$ generates the entire Hilbert space $\mathcal{H}_0$.

Although operators (L_k) are not uniquely determined for a given (τ_t) , the criteria appeared in Theorem 4.3 is independent of choice we make. We end this exposition with a comment that the result similar to Proposition 4.2 with appropriate domain condition [FR3] can be proved to include a class of quantum irreversible system where the $\mathcal{A}_0 = \mathcal{B}(\mathcal{H}_0)$ with generator which need not be a bounded operator.

REFERENCES

- [BR] Bratelli, O., Robinson, D.W. : Operator algebras and quantum statistical mechanics, I,II, Springer 1981.
- [CF] Chebotarev, A.M., Fagnola, F. Sufficient conditions for conservativity of minimal quantum dynamical semigroups, J. Funct. Anal. 153 (1998) no-2, 382-404.
- [CE] Christensen, E., Evans, D. E.: Cohomology of operator algebras and quantum dynamical semigroups, J.Lon. Maths Soc. 20 (1970) 358-368.
- [Da1] Davies, E.B.: One parameter semigroups, Academic Press, 1980.
- [Da2] Davies, E.B.: Quantum theory of open systems, Academic press, 1976.

- [Da3] Davies, E.B.: Quantum dynamical semigroups and the neutron diffusion equation. Rep. Math. Phys. vol-11, no-2 (1977), 169-188.
- [Ev] Evans, D.E.: Irreducible quantum dynamical semigroups, Commun. Math. Phys. vol-54 (1977), 293-297.
- [FR1] F. Fagnola, R. Rebolledo. The approach to equilibrium of a class of quantum dynamical semigroups Inf. Dim. Anal. Q. Prob. and Rel. Topics, vol-1, no. 4 (1998), 1–12.
- [FR2] F. Fagnola and R. Rebolledo. On the existence of invariant states for quantum dynamical semigroups J. Math. Phys., vol-42 (2001), 296-1308.
- [FR3] F. Fagnola, R. Rebolledo.: Subharmonic projections for a Quantum Markov Semigroup, J. Math. Phys., vol-43, no. 2 (2002), 1074–1082.
- [Fr] Frigerio, A.: Stationary states of quantum dynamical semigroups, Commun. Math. Phys. vol-63 (1978), 269-276.
- [GKS] Gorini, V., Kossakowski, A., Sudarshan, E.C.G. : Completely positive dynamical semigroups of n-level systems, J. Math. Phys. vol-17 (1976) 821-825.
- [Li] Lindblad, G. : On the generators of quantum dynamical semigroups, Commun. Math. Phys. vol-48 (1976), 119-130.
- [Mo1] Mohari, A.: Markov shift in Non-commutative probability, J. of Funct. Anal. vol-199, no. 1 (2003), 189–209.
- [Mo2] Mohari, A.: Ergodicity of Homogeneous Brownian flows, Stochastic processes and their application vol-105, no. 1 (2003), 99–116.

- [MS] Mohari, A., Sinha, K.B.: Stochastic dilation of minimal quantum dynamical semigroup. Proc. Indian Acad. Sci. Math Sci. vol-102 (1992), no-2, 159-173.
- [No] Norris, J.R., Markov Chain, Cambridge University Press, 1998.